

Det Kgl. Danske Videnskabernes Selskab.

Mathematisk-fysiske Meddelelser. **XIII**, 16.

THE ORBIT OF COMET 1929 I
(SCHWASSMANN-WACHMANN) IN THE
YEARS 1920—1936

BY

HANS Q. RASMUSEN

WITH 1 FIGURE



KØBENHAVN

LEVIN & MUNKSGAARD

EJNAR MUNKSGAARD

1935

Printed in Denmark.
Bianco Lunos Bogtrykkeri A/S.

The following revision of the orbit of comet 1929 I is based on the elements computed by Dr. S. KANDA (T. A. B. 41—42). These elements are as follows:

$$\begin{aligned}
 T &= 1929 \text{ March } 23.25926 \text{ U. T.} \\
 \omega &= 357.76403 \\
 \Omega &= 126.32592 \\
 i &= 3.72893 \\
 e &= 0.3950135 \\
 a &= 3.455363 \\
 \mu &= 0.1534488
 \end{aligned}
 \left. \vphantom{\begin{aligned} \omega \\ \Omega \\ i \end{aligned}} \right\} 1950.0 \tag{I}$$

$$\begin{aligned}
 x &= -1.936 \ 9388 (\cos E - e) - 2.623 \ 4602 \sin E \\
 y &= +2.628 \ 6605 (\cos E - e) - 1.710 \ 7396 \sin E \\
 z &= +1.130 \ 4612 (\cos E - e) - 0.517 \ 0711 \sin E
 \end{aligned}
 \left. \vphantom{\begin{aligned} x \\ y \end{aligned}} \right\} 1950.0$$

From these elements I have computed the perturbed co-ordinates of the comet from 1929 to 1935 (A. N. 6069 and Copenhagen Observatory Publ. 102).

The comet was rediscovered by Professor VAN BIESBROECK on 1934 Dec. 11. A comparison with the ephemeris gave the following residuals:

$$\Delta \alpha \cos \delta = +150^s, \Delta \delta = +780''.$$

From the above mentioned computation I found for 1934 Dec. 11.11181 U. T. the following approximate per-

turbations as differences between the perturbed and unperturbed co-ordinates:

$$\left. \begin{aligned} \xi &= +0.04978 \\ \eta &= -0.01415 \\ \zeta &= -0.00832 \end{aligned} \right\} 1950.0$$

From these perturbations and the following observations:¹

U. T.					
1929 Febr. 1.0	5 ^h 38 ^m 15 ^s .04	+21°10' 4".3	}	Normal place	Nr. 1 (20 obs.)
1929 March 31.0	6 35 54.50	+23 14 36.1			
1934 Dec. 11.11181	1 43 24.31	+ 5 21 31.7		Yerkes	

and using the method of the variation of the geocentric distances for the times 1929 Febr. 1.0 and 1929 March 31.0 I have computed the following revised elements:

Epoch and osculation: 1929 Febr. 18.0 U. T.

$$\left. \begin{aligned} M &= 354.90992 \\ \omega &= 357.71314 \\ \Omega &= 126.32675 \\ i &= 3.72833 \\ e &= 0.3945110 \\ a &= 3.452618 \\ \mu &= 0.1536318 \end{aligned} \right\} 1950.0 \quad (\text{II})$$

$$\left. \begin{aligned} x &= -1.932\ 906 (\cos E - e) - 2.623\ 552 \sin E \\ y &= +2.628\ 200 (\cos E - e) - 1.707\ 662 \sin E \\ z &= +1.130\ 052 (\cos E - e) - 0.515\ 907 \sin E \end{aligned} \right\} 1950.0$$

¹) All the observations and normal places are referred to the equinox for the beginning of the year of observation.

Using these elements I have repeated the computation of perturbed equatorial co-ordinates from 1929 to 1936 (7 figure computation; the perturbations from Jupiter and Saturn are considered; attention has been paid to an essential part of the effect of the planets Merkur, Venus, Earth through the use of the value $k^2 = 0.0002959139$ which corresponds to the sum of the masses of these three planets and the Sun). The co-ordinates are:

		Equinox 1950.0.					
U. T.		x		y		z	
1929	Jan.	19.0	− 0.378 379	+ 1.958 404	+ 0.782 292		
	—	29.0	0.509 634	1.915 575	0.773 037		
	Febr.	8.0	0.639 322	1.866 864	0.761 408		
	—	18.0	0.767 011	1.812 317	0.747 398		
	—	28.0	0.892 266	1.752 025	0.731 020		
	March	10.0	1.014 662	1.686 120	0.712 299		
	—	20.0	1.133 782	1.614 774	0.691 279		
	—	30.0	1.249 234	1.538 202	0.668 023		
	April	9.0	1.360 644	1.456 654	0.642 605		
	—	19.0	1.467 675	1.370 416	0.615 119		
	—	29.0	1.570 019	1.279 802	0.585 668		
	May	9.0	1.667 411	1.185 149	0.554 370		
	—	19.0	1.759 623	1.086 814	0.521 348		
	—	29.0	1.846 473	0.985 164	0.486 737		
	June	8.0	1.927 818	0.880 575	0.450 674		
	—	18.0	2.003 559	0.773 424	0.413 300		
	July	8.0	2.138 026	0.552 921	0.335 185		
	—	28.0	2.249 826	0.326 524	0.253 504		
	Aug.	17.0	2.339 409	+ 0.096 870	0.169 307		
	Sept.	6.0	2.407 610	− 0.133 698	+ 0.083 550		
	—	26.0	2.455 536	0.363 170	− 0.002 921		
	Oct.	16.0	2.484 466	0.589 859	0.089 380		
	Nov.	5.0	2.495 769	0.812 388	0.175 215		
	—	25.0	2.490 840	1.029 655	0.259 919		
	Dec.	15.0	− 2.471 056	− 1.240 804	− 0.343 081		

		Equinox 1950.0.							
U. T.		x		y		z			
1930	Jan.	4.0	— 2.437	745	— 1.445	179	— 0.424	373	
	—	24.0	2.392	165	1.642	298	0.503	536	
	Febr.	13.0	2.335	491	1.831	818	0.580	371	
	March	5.0	2.268	818	2.013	511	0.654	727	
	—	25.0	2.193	152	2.187	237	0.726	493	
	April	14.0	2.109	418	2.352	931	0.795	592	
	May	4.0	2.018	461	2.510	582	0.861	973	
	—	24.0	1.921	056	2.660	225	0.925	605	
<hr/>									
	July	3.0	1.709	660	2.935	782	1.044	583	
	Aug.	12.0	1.480	167	3.180	412	1.152	561	
	Sept.	21.0	1.236	696	3.395	151	1.249	711	
	Oct.	31.0	0.982	700	3.581	143	1.336	284	
	Dec.	10.0	0.721	095	3.739	562	1.412	571	
1931	Jan.	19.0	0.454	363	3.871	572	1.478	884	
	Febr.	28.0	— 0.184	633	3.978	295	1.535	541	
	April	9.0	+ 0.086	246	4.060	803	1.582	852	
	May	19.0	0.356	654	4.120	108	1.621	122	
	June	28.0	0.625	155	4.157	161	1.650	636	
	Aug.	7.0	0.890	461	4.172	854	1.671	672	
	Sept.	16.0	1.151	403	4.168	024	1.684	486	
	Oct.	26.0	1.406	905	4.143	453	1.689	324	
	Dec.	5.0	1.655	967	4.099	880	1.686	415	
	1932	Jan.	14.0	1.897	644	4.037	999	1.675	976
Febr.		23.0	2.131	036	3.958	464	1.658	211	
April		3.0	2.355	269	3.861	902	1.633	314	
May		13.0	2.569	489	3.748	909	1.601	467	
June		22.0	2.772	848	3.620	060	1.562	848	
Aug.		1.0	2.964	497	3.475	916	1.517	626	
Sept.		10.0	3.143	574	3.317	026	1.465	966	
Oct.		20.0	3.309	196	3.143	936	1.408	030	
Nov.		29.0	3.460	452	2.957	196	1.343	981	
1933		Jan.	8.0	3.596	392	2.757	367	1.273	983
	Febr.	17.0	3.716	016	2.545	031	1.198	206	
	March	29.0	3.818	266	2.320	800	1.116	828	
	May	8.0	+ 3.902	018	— 2.085	329	— 1.030	041	

		Equinox 1950.0.						
U. T.		x		y		z		
1933	June	17.0	+ 3.966	066	- 1.839	335	- 0.938	055
	July	27.0	4.009	110	1.583	610	0.841	105
	Sept.	5.0	4.029	748	1.319	045	0.739	457
	Oct.	15.0	4.026	457	1.046	662	0.633	424
	Nov.	24.0	3.997	585	0.767	647	0.523	373
1934	Jan.	3.0	3.941	330	0.483	397	0.409	746
	Febr.	12.0	3.855	738	- 0.195	576	0.293	078
	March	24.0	3.738	689	+ 0.093	806	0.174	033
	May	3.0	3.587	908	0.382	297	- 0.053	435
	June	12.0	3.400	977	0.666	878	+ 0.067	680
	July	2.0	3.293	176	0.806	566	0.128	033
	—	22.0	3.175	397	0.943	805	0.187	997
	Aug.	11.0	3.047	332	1.077	981	0.247	349
	—	31.0	2.908	684	1.208	406	0.305	840
	Sept.	20.0	2.759	176	1.334	310	0.363	185
	Oct.	10.0	2.598	564	1.454	836	0.419	065
	—	30.0	2.426	653	1.569	022	0.473	118
	Nov.	19.0	2.243	310	1.675	802	0.524	937
	Dec.	9.0	2.048	496	1.773	996	0.574	065
	—	29.0	1.842	288	1.862	301	0.619	992
1935	Jan.	18.0	1.624	919	1.939	298	0.662	153
	Febr.	7.0	1.396	820	2.003	455	0.699	928
	—	27.0	1.158	666	2.053	147	0.732	648
	March	19.0	0.911	427	2.086	688	0.759	604
	April	8.0	0.656	422	2.102	377	0.780	058
	—	28.0	0.395	363	2.098	570	0.793	278
	May	18.0	+ 0.130	383	2.073	775	0.798	562
	June	7.0	- 0.135	962	2.026	760	0.795	289
	—	27.0	0.400	729	1.956	679	0.782	962
	July	17.0	0.660	677	1.863	191	0.761	268
	Aug.	6.0	0.912	399	1.746	560	0.730	117
	—	26.0	1.152	498	1.607	708	0.689	676
	Sept.	15.0	1.377	792	1.448	207	0.640	378
Oct.	5.0	1.585	496	1.270	207	0.582	899	
—	25.0	- 1.773	376	+ 1.076	307	+ 0.518	126	

From these co-ordinates α and δ have been found through interpolation and reduction to the equinox for the beginning of the year of observation, and the following residuals were found:

U. T.		$\Delta \alpha \cos \delta$	$\Delta \delta$		
1934 Dec.	11.11181	— 12 ^s 73	— 65 ^{''} 2	Yerkes	
—	—	11.17500	— 12.87	— 67.2	»
—	—	12.10558	— 12.75	— 65.9	»
1935 Febr.	6.80389	— 11.01	— 60.9	Bergedorf	
—	—	6.84194	— 12.03	— 67.1	»
—	March	5.80424	— 11.64	— 62.5	»

which led to the normal places:

U. T.				
1934 Dec.	12.0	1 ^h 43 ^m 14 ^s 64	+ 5°22' 9 ^{''} 6	Normal place Nr. 3
1935 Febr.	7.0	2 11 57.78	+ 9 37 36.1	— — - 4

Comparison with all the normal places gave the result:

	U. T.		$\Delta \alpha \cos \delta$	$\Delta \delta$	
Normal place Nr. 1	1929 Febr.	1.0	— 0 ^s 20	— 0 ^{''} 5	
—	—	- 2 1929 March	31.0	+ 0.17	— 0.2
—	—	- 3 1934 Dec.	12.0	— 12.79	— 66.1
—	—	- 4 1935 Febr.	7.0	— 11.43	— 62.5

As the large residuals in 1934 and 1935 presumably arise from the inaccuracy of the perturbations used for the revision I have computed new perturbations for 1934 Dec. 11 from the difference between the new perturbed co-ordinates and the unperturbed co-ordinates deduced from the elements (II). The new perturbations are for 1934 Dec. 11.11181 U. T.:

$$\left. \begin{aligned} \xi &= +0.048\ 392 \\ \eta &= -0.013\ 344 \\ \zeta &= -0.007\ 886 \end{aligned} \right\} 1950.0$$

From these values of the perturbations I have made a new revision, which gave the new set of elements:

Epoch and osculation 1929 Febr. 18.0 U. T.

$$\left. \begin{aligned} M &= 354^{\circ}91062 \\ \omega &= 357.71203 \\ \Omega &= 126.32728 \\ i &= 3.72839 \\ e &= 0.3945470 \\ a &= 3.452821 \\ \mu &= 0.1536183 \end{aligned} \right\} 1950.0 \quad \text{(III)}$$

$$\left. \begin{aligned} x &= -1.932\ 993 (\cos E - e) - 2.623\ 676 \sin E \\ y &= +2.628\ 374 (\cos E - e) - 1.707\ 710 \sin E \\ z &= +1.130\ 122 (\cos E - e) - 0.515\ 915 \sin E \end{aligned} \right\} 1950.0$$

Using these elements the computation of the perturbed equatorial co-ordinates for 1929—1936 has once more been repeated with the following result:

		Equinox 1950.0.					
U.T.		x		y		z	
1929	Jan. 19.0	-0.378	371	+1.958	408	+0.782	290
	— 29.0	0.509	628	1.915	578	0.773	035
	Febr. 8.0	0.639	318	1.866	866	0.761	406
	— 18.0	0.767	008	1.812	318	0.747	397
	— 28.0	0.892	265	1.752	026	0.731	018
	March 10.0	1.014	662	1.686	119	0.712	297
	— 20.0	1.133	784	1.614	772	0.691	277
	— 30.0	-1.249	237	+1.538	199	+0.668	020

		Equinox 1950.0.					
U. T.		x		y		z	
1929	April	9.0	-1.360 649	+1.456 650	+0.642 603		
	—	19.0	1.467 681	1.370 412	0.615 117		
	—	29.0	1.570 027	1.279 797	0.585 666		
	May	9.0	1.667 420	1.185 143	0.554 367		
	—	19.0	1.759 635	1.086 806	0.521 346		
	—	29.0	1.846 486	0.985 155	0.486 734		
	June	8.0	1.927 833	0.880 566	0.450 672		
	—	18.0	2.003 576	0.773 414	0.413 298		
	July	8.0	2.138 045	0.552 909	0.335 183		
	—	28.0	2.249 848	0.326 511	0.253 502		
	Aug.	17.0	2.339 435	+0.096 857	0.169 305		
	Sept.	6.0	2.407 641	-0.133 713	+0.083 549		
	—	26.0	2.455 572	0.363 186	-0.002 922		
	Oct.	16.0	2.484 507	0.589 876	0.089 381		
	Nov.	5.0	2.495 816	0.812 405	0.175 215		
	—	25.0	2.490 894	1.029 674	0.259 919		
	Dec.	15.0	2.471 118	1.240 824	0.343 081		
1930	Jan.	4.0	2.437 815	1.445 201	0.424 372		
	—	24.0	2.392 242	1.642 322	0.503 535		
	Febr.	13.0	2.335 578	1.831 845	0.580 370		
	March	5.0	2.268 914	2.013 541	0.654 726		
	—	25.0	2.193 258	2.187 271	0.726 493		
	April	14.0	2.109 533	2.352 969	0.795 594		
	May	4.0	2.018 588	2.510 625	0.861 976		
	—	24.0	1.921 193	2.660 274	0.925 609		
	July	3.0	1.709 819	2.935 845	1.044 590		
	Aug.	12.0	1.480 348	3.180 491	1.152 574		
	Sept.	21.0	1.236 899	3.395 249	1.249 730		
	Oct.	31.0	0.982 925	3.581 263	1.336 310		
	Dec.	10.0	0.721 341	3.739 706	1.412 605		
1931	Jan.	19.0	0.454 628	3.871 749	1.478 927		
	Febr.	28.0	-0.184 917	3.978 497	1.535 594		
	April	9.0	+0.085 945	4.061 037	1.582 918		
	May	19.0	0.356 337	4.120 377	1.621 200		
	June	28.0	+0.624 825	-4.157 467	-1.650 728		

		Equinox 1950.0.					
U. T.		x		y		z	
1931	Aug.	7.0	+ 0.890 120	- 4.173 199	- 1.671 779		
	Sept.	16.0	1.151 053	4.168 410	1.684 609		
	Oct.	26.0	1.406 549	4.143 884	1.689 464		
	Dec.	5.0	1.655 608	4.100 356	1.686 573		
1932	Jan.	14.0	1.897 286	4.038 522	1.676 153		
	Febr.	23.0	2.130 682	3.959 037	1.658 408		
	April	3.0	2.354 923	3.862 525	1.633 531		
	May	13.0	2.569 154	3.749 584	1.601 706		
	June	22.0	2.772 530	3.620 789	1.563 109		
	Aug.	1.0	2.964 199	3.476 700	1.517 910		
	Sept.	10.0	3.143 301	3.317 865	1.466 274		
	Oct.	20.0	3.308 955	3.144 832	1.408 363		
	Nov.	29.0	3.460 247	2.958 150	1.344 339		
1933	Jan.	8.0	3.596 230	2.758 378	1.274 366		
	Febr.	17.0	3.715 903	2.546 099	1.198 615		
	March	29.0	3.818 210	2.321 925	1.117 264		
	May	8.0	3.902 027	2.086 511	1.030 503		
	June	17.0	3.966 147	1.840 572	0.938 543		
	July	27.0	4.009 273	1.584 899	0.841 619		
	Sept.	5.0	4.030 003	1.320 384	0.739 997		
	Oct.	15.0	4.026 816	1.048 048	0.633 988		
	Nov.	24.0	3.998 059	0.769 074	0.523 961		
1934	Jan.	3.0	3.941 932	0.484 857	0.410 354		
	Febr.	12.0	3.856 482	- 0.197 062	0.293 705		
	March	24.0	3.739 590	+ 0.092 306	0.174 675		
	May	3.0	3.588 981	0.380 796	- 0.054 087		
	June	12.0	3.402 240	0.665 395	+ 0.067 024		
	July	2.0	3.294 540	0.805 100	0.127 378		
	—	22.0	3.176 867	0.942 363	0.187 345		
	Aug.	11.0	3.048 912	1.076 570	0.246 703		
	—	31.0	2.910 377	1.207 033	0.305 202		
	Sept.	20.0	2.760 986	1.332 986	0.362 560		
	Oct.	10.0	2.600 496	1.453 570	0.418 456		
	—	30.0	2.428 707	1.567 827	0.472 530		
	Nov.	19.0	+ 2.245 490	+ 1.674 692	+ 0.524 375		

Equinox 1950.0.

U. T.		x	y	z
1934	Dec. 9.0	+2.050 800	+1.772 983	+0.573 534
	— 29.0	1.844 716	1.861 401	0.619 498
1935	Jan. 18.0	1.627 468	1.938 529	0.661 703
	Febr. 7.0	1.399 484	2.002 835	0.699 531
	— 27.0	1.161 436	2.052 696	0.732 311
	March 19.0	0.914 290	2.086 424	0.759 335
	April 8.0	0.659 362	2.102 320	0.779 866
	— 28.0	0.398 358	2.098 739	0.793 171
	May 18.0	+0.133 406	2.074 185	0.798 548
	June 7.0	-0.132 942	2.027 424	0.795 374
	— 27.0	0.397 748	1.957 603	0.783 152
	July 17.0	0.657 774	1.864 377	0.761 565
	Aug. 6.0	0.909 609	1.748 001	0.730 520
	— 26.0	1.149 862	1.609 389	0.690 182
	Sept. 15.0	1.375 343	1.450 106	0.640 980
	Oct. 5.0	1.583 262	1.272 298	0.583 589
	— 25.0	1.771 378	1.078 558	0.518 892
	Nov. 14.0	1.938 090	0.871 761	0.447 915
	Dec. 4.0	2.082 459	0.654 885	0.371 764
	— 24.0	2.204 172	0.430 856	0.291 556
1936	Jan. 13.0	-2.303 859	+0.202 413	+0.208 371

Comparison with the 4 normal places then gave the residuals:

U. T.		$\Delta a \cos \delta$	$\Delta \delta$
Normal place Nr. 1	1929 Febr. 1.0	-0 ^s .15	+0 ^{''} .2
— — - 2	1929 March 31.0	+0.14	+0.1
— — - 3	1934 Dec. 12.0	+0.02	+2.3
— — - 4	1935 Febr. 7.0	-0.89	-5.5

The discrepancies between observation and computation are now so small that the orbit may be considered as fairly satisfactory. A revision based on the last computation of

perturbed co-ordinates should be postponed till the appearance of new observations.

From the co-ordinates and velocities for 1935 Oct. 5.0 I deduced the following osculating elements:

Epoch and osculation 1935 Oct. 5.0 U. T.

$$\left. \begin{aligned} M &= 5.74536 \\ \omega &= 357.97151 \\ \Omega &= 126.29158 \\ i &= 3.72853 \end{aligned} \right\} 1950.0 \quad \text{(III a)}$$

$$\begin{aligned} e &= 0.3938329 \\ a &= 3.455860 \\ \mu &= 0.1534157 \end{aligned}$$

$$\left. \begin{aligned} x &= -1.945\ 836 (\cos E - e) - 2.619\ 892 \sin E \\ y &= +2.623\ 348 (\cos E - e) - 1.719\ 222 \sin E \\ z &= +1.129\ 044 (\cos E - e) - 0.520\ 587 \sin E \end{aligned} \right\} 1950.0$$

As it may be possible to observe the comet during the winter 1935—36 I have computed the following ephemeris from the elements (III a).

0^h U. T.

1935		α 1935.0	δ 1935.0	Δ	r
Oct.	25	11 ^h 6 ^m 49 ^s	+ 6°46'1	2.714	2.138
—	29	14 33	6 3.0	2.687	2.144
Nov.	2	22 10	5 20.3	2.659	2.150
—	6	29 39	4 38.1	2.631	2.157
—	10	37 1	3 56.5	2.602	2.164
—	14	44 16	3 15.6	2.572	2.172
—	18	51 23	2 35.5	2.541	2.180
—	22	11 58 21	1 56.4	2.509	2.188
—	26	12 5 10	1 18.4	2.477	2.196
—	30	12 11 49	+ 0 41.6	2.444	2.205

0^h U. T.

	1935	α 1935.0	δ 1935.0	Δ	r
Dec.	4	12 ^h 18 ^m 19 ^s	+ 0° 6′.0	2.410	2.214
—	8	24 39	— 0 28.1	2.375	2.224
—	12	30 48	1 0.7	2.340	2.234
—	16	36 45	1 31.8	2.305	2.244
—	20	42 30	2 1.2	2.269	2.254
—	24	48 1	2 28.7	2.232	2.265
—	28	53 18	2 54.3	2.195	2.276
—	32	12 58 21	— 3 17.9	2.158	2.287

0^h U. T.

	1936	α 1936.0	δ 1936.0	Δ	r
Jan.	1	12 ^h 58 ^m 24 ^s	— 3°18′.2	2.158	2.287
—	5	13 3 11	3 39.8	2.120	2.298
—	9	7 40	3 59.1	2.083	2.310
—	13	11 52	4 16.2	2.045	2.322
—	17	15 44	4 30.9	2.008	2.334
—	21	19 16	4 43.2	1.971	2.346
—	25	22 27	4 53.0	1.934	2.359
—	29	25 15	5 0.2	1.898	2.371
Febr.	2	27 39	5 4.9	1.863	2.384
—	6	29 38	5 6.9	1.830	2.397
—	10	31 12	5 6.3	1.797	2.411
—	14	32 20	5 3.0	1.766	2.424
—	18	33 1	4 57.1	1.736	2.438
—	22	33 15	4 48.6	1.709	2.452
—	26	33 1	4 37.7	1.683	2.466
March	1	32 21	4 24.5	1.660	2.480
—	5	13 31 15	— 4 9.0	1.641	2.494

The small inclination of the orbit and the distance of the aphelion 4.81 made a former proximity to Jupiter probable. In fact such a proximity took place in the first part of 1926.

To find the elements before this I have computed the co-ordinates backwards till 1920. From 1929 to 1924 Jupiter and Saturn were taken into consideration in a 7-figure computation, from 1924 to 1920 only Jupiter was considered and a 6-figure computation was used. The resulting equatorial co-ordinates were as follows:

		Equinox 1950.0.					
U. T.		x		y		z	
1929	Jan.	9.0	- 0.245 974	+ 1.995 348	+ 0.789 193		
1928	Dec.	30.0	- 0.112 853	2.026 431	0.793 779		
	—	20.0	+ 0.020 594	2.051 722	0.796 096		
	—	10.0	0.153 986	2.071 320	0.796 205		
	Nov.	30.0	0.286 968	2.085 352	0.794 173		
	—	20.0	0.419 207	2.093 966	0.790 078		
	—	10.0	0.550 399	2.097 332	0.784 002		
	Oct.	21.0	0.808 557	2.089 076	0.766 266		
	—	1.0	1.059 544	2.062 193	0.741 695		
	Sept.	11.0	1.301 874	2.018 388	0.711 037		
	Aug.	22.0	1.534 436	1.959 377	0.675 020		
	—	2.0	1.756 437	1.886 832	0.634 340		
	July	13.0	1.967 356	1.802 341	0.589 642		
	June	23.0	2.166 890	1.707 383	0.541 520		
	—	3.0	2.354 908	1.603 320	0.490 508		
	May	14.0	2.531 414	1.491 391	0.437 091		
	April	24.0	2.696 514	1.372 718	0.381 696		
	—	4.0	2.850 391	1.248 308	0.324 707		
	March	15.0	2.993 281	1.119 066	0.266 461		
	Febr.	24.0	3.125 459	0.985 800	0.207 258		
	—	4.0	3.247 225	0.849 234	0.147 362		
	Jan.	15.0	3.358 894	0.710 012	0.087 008		
1927	Dec.	26.0	3.460 791	0.568 712	+ 0.026 400		
	—	6.0	3.553 241	0.425 849	- 0.034 276		
	Nov.	16.0	3.636 570	0.281 885	0.094 860		
	Oct.	27.0	3.711 097	+ 0.137 234	0.155 209		
	—	7.0	+ 3.777 136	- 0.007 734	- 0.215 194		

Equinox 1950.0.

	U. T.		x		y		z
1927	Sept.	17.0	+ 3.834	994	- 0.152	682	- 0.274 701
	Aug.	28.0	3.884	966	0.297	311	0.333 629
	—	8.0	3.927	341	0.441	348	0.391 887
	July	19.0	3.962	396	0.584	546	0.449 394
	June	29.0	3.990	401	0.726	680	0.506 076
	—	9.0	4.011	615	0.867	547	0.561 866
	May	20.0	4.026	290	1.006	962	0.616 706
	April	30.0	4.034	668	1.144	752	0.670 540
	—	10.0	4.036	986	1.280	763	0.723 319
	March	21.0	4.033	473	1.414	852	0.774 999
	—	1.0	4.024	353	1.546	885	0.825 538
	Febr.	9.0	4.009	845	1.676	742	0.874 899
	Jan.	20.0	3.990	164	1.804	308	0.923 048
1926	Dec.	31.0	3.965	524	1.929	483	0.969 954
	—	11.0	3.936	138	2.052	170	1.015 591
	Nov.	21.0	3.902	221	2.172	283	1.059 933
	—	1.0	3.863	992	2.289	745	1.102 963
	Oct.	12.0	3.821	679	2.404	491	1.144 665
	Sept.	22.0	3.775	517	2.516	467	1.185 032
	—	2.0	3.725	760	2.625	634	1.224 067
	Aug.	13.0	3.672	683	2.731	979	1.261 787
	July	24.0	3.616	584	2.835	523	1.298 233
	—	4.0	3.557	790	2.936	335	1.333 481
	June	14.0	3.496	654	3.034	561	1.367 658
	May	25.0	3.433	521	3.130	453	1.400 974
	—	5.0	3.368	666	3.224	398	1.433 737
	April	15.0	3.302	173	3.316	904	1.466 350
	March	26.0	3.233	796	3.408	483	1.499 229
	—	6.0	3.162	945	3.499	449	1.532 652
	Febr.	14.0	3.088	901	3.589	738	1.566 614
	Jan.	25.0	3.011	102	3.678	972	1.600 869
	—	5.0	2.929	293	3.766	630	1.635 054
1925	Dec.	16.0	2.843	474	3.852	210	1.668 808
	Nov.	26.0	2.753	801	3.935	314	1.701 847
	—	6.0	+ 2.660	495	- 4.015	635	- 1.733 945

		Equinox 1950.0.					
U. T. (resp. G. M. T.)		x		y		z	
1925	Oct.	17.0	+ 2.563 797	- 4.092 950	- 1.764 944		
	Sept.	27.0	2.463 944	4.167 095	1.794 725		
	—	7.0	2.361 157	4.237 948	1.823 204		
	Aug.	18.0	2.255 642	4.305 413	1.850 313		
	July	29.0	2.147 592	4.369 419	1.876 007		
	—	9.0	2.037 184	4.429 907	1.900 244		
	June	19.0	1.924 583	4.486 828	1.922 997		
	May	10.0	1.693 423	4.589 815	1.963 950		
	March	31.0	1.455 270	4.678 125	1.998 718		
	Febr.	19.0	1.211 194	4.751 553	2.027 193		
	Jan.	10.0	0.962 186	4.809 941	2.049 293		
1924	Nov.	30.5	0.709 202	4.853 159	2.064 958		
	Oct.	21.5	0.453 158	4.881 090	2.074 138		
	Sept.	11.5	+ 0.194 951	4.893 642	2.076 794		
	Aug.	2.5	- 0.064 537	4.890 738	2.072 895		
	June	23.5	0.324 425	4.872 310	2.062 418		
	May	14.5	0.583 839	4.838 312	2.045 350		
	April	4.5	0.841 895	4.788 711	2.021 681		
	Febr.	24.5	1.097 700	4.723 492	1.991 412		
	Jan.	15.5	1.350 351	4.642 655	1.954 551		
1923	Dec.	6.5	1.598 922	4.546 228	1.911 117		
	Oct.	27.5	1.842 469	4.434 256	1.861 139		
	Sept.	17.5	2.080 022	4.306 816	1.804 657		
	Aug.	8.5	2.310 586	4.164 013	1.741 723		
	June	29.5	2.533 131	4.005 993	1.672 410		
	May	20.5	2.746 594	3.832 938	1.596 803		
	April	10.5	2.949 880	3.645 084	1.515 015		
	March	1.5	3.141 853	3.442 716	1.427 176		
	Jan.	20.5	3.321 341	3.226 190	1.333 448		
1922	Dec.	11.5	3.487 132	2.995 929	1.234 023		
	Nov.	1.5	3.637 975	2.752 445	1.129 128		
	Sept.	22.5	3.772 587	2.496 342	1.019 034		
	Aug.	13.5	3.889 646	2.228 334	0.904 055		
	July	4.5	- 3.987 810	- 1.949 257	- 0.784 560		

Equinox 1950.0.

G. M. T.		x	y	z
1922	May 25.5	-4.065 714	-1.660 081	-0.660 977
	April 15.5	4.121 987	1.361 927	0.533 793
	March 6.5	4.155 273	1.056 079	0.403 571
	Jan. 25.5	4.164 239	0.743 996	0.270 945
1921	Dec. 16.5	4.147 611	0.427 329	0.136 633
	Nov. 6.5	4.104 207	-0.107 925	-0.001 433
	Sept. 27.5	4.032 966	+0.212 161	+0.133 769
	Aug. 18.5	3.933 002	0.530 666	0.268 000
	July 9.5	3.803 644	0.845 121	0.400 201
	May 30.5	3.644 491	1.152 864	0.529 235
	April 20.5	3.455 468	1.451 060	0.653 897
	March 11.5	3.236 873	1.736 738	0.772 925
	Jan. 30.5	2.989 433	2.006 841	0.885 026
1920	Dec. 21.5	-2.714 342	+2.258 280	+0.988 902

The figure on page 19 illustrates the abnormally large perturbations in the orbit of the comet near the proximity to Jupiter in 1926. The orbit is shown projected on the ecliptic (the inclination is always very small).


From the co-ordinates and velocities for 1921 March 11.5 I have deduced the following set of elements:

Epoch and osculation 1921 March 11.5 G. M. T.

$$\begin{aligned}
 M &= 32.384 \\
 \omega &= 335.462 \\
 \Omega &= 126.681 \\
 i &= 0.710 \\
 e &= 0.19707 \\
 a &= 4.4261 \\
 \mu &= 0.10585
 \end{aligned}
 \left. \vphantom{\begin{aligned} M \\ \omega \\ \Omega \\ i \\ e \\ a \\ \mu \end{aligned}} \right\} 1950.0 \quad \text{(III b)}$$

$$\begin{aligned}
 x &= -0.93118 (\cos E - e) - 4.24194 \sin E \\
 y &= +3.97877 (\cos E - e) - 0.85674 \sin E \\
 z &= +1.70072 (\cos E - e) - 0.31824 \sin E
 \end{aligned}
 \left. \vphantom{\begin{aligned} x \\ y \\ z \end{aligned}} \right\} 1950.0$$

I have not been successful in establishing any identity between these elements and any other set of elements in the tables of comets and asteroids.



The following table shows the distance of the comet from Jupiter for a number of dates from 1921 to 1928:

U. T. (resp. G. M. T.)	Δ
1928 Nov. 10.0	3.508
1927 Oct. 7.0	1.188
1926 Sept. 2.0	0.357
1925 July 29.0	0.438
1924 June 23.5	0.895
1923 May 20.5	1.078
1922 April 15.5	1.141
1921 March 11.5	2.055

On 1926 March 26 the distance had a minimum value:
0.179.

I am indebted to the Carlsbergfond for a grant, which
enabled me to carry out the present investigation.

University Observatory, Copenhagen 1935 Aug. 9.

HANS Q. RASMUSEN.